

On n -maximal subgroups of finite groups

Vika A. Kovaleva

Department of Mathematics, Francisk Skorina Gomel State University,
Gomel 246019, Belarus
E-mail: vika.kovalyova@rambler.ru

Alexander N. Skiba

Department of Mathematics, Francisk Skorina Gomel State University,
Gomel 246019, Belarus
E-mail: alexander.skiba49@gmail.com

Abstract

We describe finite soluble groups in which every n -maximal subgroup is $\mathcal{F}$ -subnormal for some saturated formation $\mathcal{F}$.

1 Introduction

Throughout this paper, all groups are finite and G always denotes a finite group. We use $\mathcal{U}$, $\mathcal{N}$ and $\mathcal{N}^r$ to denote the class of all supersoluble groups, the class of all nilpotent groups and the class of soluble groups of nilpotent length at most r ($r \geq 1$). The symbol $\mathbb{P}$ denotes the set of all primes, $\pi(G)$ denotes the set of prime divisors of the order of G . If p is a prime, then we use $\mathcal{G}_p$ to denote the class of all p -groups.

Let $\mathcal{F}$ be a class of groups. If $1 \in \mathcal{F}$, then we write $G^{\mathcal{F}}$ to denote the intersection of all normal subgroups N of G with $G/N \in \mathcal{F}$. The class $\mathcal{F}$ is said to be a *formation* if either $\mathcal{F} = \emptyset$ or $1 \in \mathcal{F}$ and every homomorphic image of $G/G^{\mathcal{F}}$ belongs to $\mathcal{F}$ for any group G . The formation $\mathcal{F}$ is said to be: *saturated* if $G \in \mathcal{F}$ whenever $G/\Phi(G) \in \mathcal{F}$ for any group G ; *hereditary* if $H \in \mathcal{F}$ whenever $G \in \mathcal{F}$ and H is a subgroup of G . A group G is called $\mathcal{F}$ -critical provided G does not belong to $\mathcal{F}$ but every proper subgroup of G belongs to $\mathcal{F}$.

For any formation function $f : \mathbb{P} \rightarrow \{\text{group formation}\}$, the symbol $LF(f)$ denotes the collection of all groups G such that either $G = 1$ or $G \neq 1$ and $G/C_G(H/K) \in f(p)$ for every chief factor H/K of G and every $p \in \pi(H/K)$. It is well-known that for any non-empty saturated formation $\mathcal{F}$, there

Keywords: n -maximal subgroup, soluble group, supersoluble group, n -multiply saturated formation, $\mathcal{F}$ -critical group, $\mathcal{F}$ -subnormal subgroup.

Mathematics Subject Classification (2000): 20D10, 20D15

is a unique formation function F such that $\mathcal{F} = LF(F)$ and $F(p) = \mathcal{G}_p F(p) \subseteq \mathcal{F}$ for all primes p , where $\mathcal{G}_p F(p)$ is the set of all groups G such that $G^{F(p)} \in \mathcal{G}_p$ (see Proposition 3.8 in [1, Chapter IV]). The formation function F is called the *canonical local satellite* of $\mathcal{F}$. A chief factor H/K of G is called $\mathcal{F}$ -*central* in G provided $G/C_G(H/K) \in F(p)$ for all primes p dividing $|H/K|$, otherwise it is called $\mathcal{F}$ -*eccentric*.

Fix some ordering ϕ of $\mathbb{P}$. The record $p\phi q$ means that p precedes q in ϕ and $p \neq q$. Recall that a group G of order $p_1^{\alpha_1} p_2^{\alpha_2} \dots p_n^{\alpha_n}$ is called ϕ -*dispersive* whenever $p_1 \phi p_2 \phi \dots \phi p_n$ and for every i there is a normal subgroup of G of order $p_1^{\alpha_1} p_2^{\alpha_2} \dots p_i^{\alpha_i}$. Furthermore, if ϕ is such that $p\phi q$ always implies $p > q$, then every ϕ -dispersive group is called *Ore dispersive*.

By definition, every formation is *0-multiply saturated* and for $n \geq 1$ a formation $\mathcal{F}$ is called *n-multiply saturated* if $\mathcal{F} = LF(f)$, where every non-empty value of the function f is an $(n-1)$ -multiply saturated formation (see [2] and [3]). In fact, almost saturated formations met in mathematical practice are n -multiply saturated for every natural n . For example, the formations of all soluble groups, all nilpotent groups, all p -soluble groups, all p -nilpotent groups, all p -closed groups, all p -decomposable groups, all Ore dispersive groups, all metanilpotent groups are n -multiply saturated for all $n \geq 1$. Nevertheless, the formations of all supersoluble groups and all p -supersoluble groups are saturated, but they are not 2-multiply saturated formations.

Recall that a subgroup H of G is called a *2-maximal (second maximal)* subgroup of G whenever H is a maximal subgroup of some maximal subgroup M of G . Similarly we can define *3-maximal subgroups*, and so on.

The interesting and substantial direction in finite group theory consists in studying the relations between the structure of the group and its n -maximal subgroups. One of the earliest publication in this direction is the article of Huppert [4] who established the supersolubility of a group G whose all second maximal subgroups are normal. In the same article Huppert proved that if all 3-maximal subgroups of G are normal in G , then the commutator subgroup G' of G is nilpotent and the chief rank of G is at most 2. These two results were developed by many authors. Among the recent results on n -maximal subgroups we can mention [5], where the solubility of groups is established in which all 2-maximal subgroups enjoy the cover-avoidance property, and [6, 7, 8], where new characterizations of supersoluble groups in terms of 2-maximal subgroups were obtained. The classification of non-nilpotent groups whose all 2-maximal subgroups are *TI*-subgroups appeared in [9]. Description was obtained in [10] of groups whose every 3-maximal subgroup permutes with all maximal subgroups. The nonnilpotent groups are described in [11] in which every two 3-maximal subgroups are permutable. The groups are described in [12] whose all 3-maximal subgroups are *S*-quasinormal, that is, permute with all Sylow subgroups. Subsequently this result was strengthened in [13] to provide a description of the groups whose all 3-maximal subgroups are subnormal.

Despite of all these and many other known results about n -maximal subgroups, the fundamental work of Mann [14] still retains its value. It studied the structure of groups whose n -maximal

subgroups are subnormal. Mann proved that if all n -maximal subgroups of a soluble group G are subnormal and $|\pi(G)| \geq n + 1$, then G is nilpotent; but if $|\pi(G)| \geq n - 1$, then G is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$. Finally, in the case $|\pi(G)| = n$ Mann described G completely.

Let $\mathcal{F}$ be a non-empty formation. Recall that a subgroup H of a group G is said to be $\mathcal{F}$ -subnormal in G if either $H = G$ or there exists a chain of subgroups $H = H_0 < H_1 < \dots < H_n = G$ such that H_{i-1} is a maximal subgroup of H_i and $H_i/(H_{i-1})_{H_i} \in \mathcal{F}$, for $i = 1, \dots, n$.

The main goal of this article is to prove the following formation analogs of Mann's theorems.

Theorem A. *Let $\mathcal{F}$ be an r -multiply saturated formation such that $\mathcal{N} \subseteq \mathcal{F} \subseteq \mathcal{N}^{r+1}$ for some $r \geq 0$. If every n -maximal subgroup of a soluble group G is $\mathcal{F}$ -subnormal in G and $|\pi(G)| \geq n + r + 1$, then $G \in \mathcal{F}$.*

Theorem B. *Let $\mathcal{F} = LF(F)$ be a saturated formation such that $\mathcal{N} \subseteq \mathcal{F} \subseteq \mathcal{U}$, where F is the canonical local satellite of $\mathcal{F}$. Let G be a soluble group with $|\pi(G)| \geq n + 1$. Then all n -maximal subgroups of G are $\mathcal{F}$ -subnormal in G if and only if G is a group of one of the following types:*

I. $G \in \mathcal{F}$.

II. $G = A \rtimes B$, where $A = G^{\mathcal{F}}$ and B are Hall subgroups of G , while G is Ore dispersive and satisfies the following:

(1) A is either of the form $N_1 \times \dots \times N_t$, where each N_i is a minimal normal subgroup of G , which is a Sylow subgroup of G , for $i = 1, \dots, t$, or a Sylow p -subgroup of G of exponent p for some prime p and the commutator subgroup, the Frattini subgroup, and the center of A coincide, while $A/\Phi(A)$ is an $\mathcal{F}$ -eccentric chief factor of G ;

(2) every n -maximal subgroup of G belongs to $\mathcal{F}$ and induces on the Sylow p -subgroup of A an automorphism group which is contained in $F(p)$ for every prime divisor p of $|A|$.

In the proof of Theorem B we often use Theorem A and the following useful fact.

Theorem C. *Let $\mathcal{F}$ be a hereditary saturated formation such that every $\mathcal{F}$ -critical group is soluble and it has a normal Sylow p -subgroup $G_p \neq 1$ for some prime p . Then every 2-maximal subgroup of G is $\mathcal{F}$ -subnormal in G if and only if either $G \in \mathcal{F}$ or G is an $\mathcal{F}$ -critical group and $G^{\mathcal{F}}$ is a minimal normal subgroup of G .*

Theorem D. *Let $\mathcal{F}$ be a saturated formation such that $\mathcal{N} \subseteq \mathcal{F} \subseteq \mathcal{U}$. If every n -maximal subgroup of a soluble group G is $\mathcal{F}$ -subnormal in G and $|\pi(G)| \geq n$, then G is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$.*

All unexplained notation and terminology are standard. The reader is referred to [1] or [15] if necessary.

2 Preliminary Results

Let $\mathcal{F}$ be a non-empty formation. Recall that a maximal subgroup H of G is said to be $\mathcal{F}$ -normal in G if $G/H_G \in \mathcal{F}$, otherwise it is said to be $\mathcal{F}$ -abnormal in G .

We use the following results.

Lemma 2.1. *Let $\mathcal{F}$ be a formation and H an $\mathcal{F}$ -subnormal subgroup of G .*

(1) *If $\mathcal{F}$ is hereditary and $K \leq G$, then $H \cap K$ is an $\mathcal{F}$ -subnormal subgroup in K [15, Lemma 6.1.7(2)].*

(2) *If N is a normal subgroup in G , then HN/N is an $\mathcal{F}$ -subnormal subgroup in G/N [15, Lemma 6.1.6(3)].*

(3) *If K is a subgroup of G such that K is $\mathcal{F}$ -subnormal in H , then K is $\mathcal{F}$ -subnormal in G [15, Lemma 6.1.6(1)].*

(4) *If $\mathcal{F}$ is hereditary and K is a subgroup of G such that $G^{\mathcal{F}} \leq K$, then K is $\mathcal{F}$ -subnormal in G [15, Lemma 6.1.7(1)].*

The following lemma is evident.

Lemma 2.2. *Let $\mathcal{F}$ be a hereditary formation. If $G \in \mathcal{F}$, then every subgroup of G is $\mathcal{F}$ -subnormal in G .*

Lemma 2.3. *Let $\mathcal{F}$ be a hereditary saturated formation. If every n -maximal subgroup of G is $\mathcal{F}$ -subnormal in G , then every $(n-1)$ -maximal subgroup of G belongs to $\mathcal{F}$ and every $(n+1)$ -maximal subgroup of G is $\mathcal{F}$ -subnormal in G .*

Proof. We first show that every $(n-1)$ -maximal subgroup of G belongs to $\mathcal{F}$. Let H be an $(n-1)$ -maximal subgroup of G and K a maximal subgroup of H . Then K is an n -maximal subgroup of G and so, by hypothesis, K is $\mathcal{F}$ -subnormal in G . Hence K is $\mathcal{F}$ -subnormal in H by Lemma 2.1(1). Thus all maximal subgroups of H are $\mathcal{F}$ -normal in H . Therefore $H \in \mathcal{F}$ since $\mathcal{F}$ is saturated.

Now, let E be an $(n+1)$ -maximal subgroup of G , and let E_1 and E_2 be an n -maximal and an $(n-1)$ -maximal subgroup of G , respectively, such that $E \leq E_1 \leq E_2$. Then, by the above, $E_2 \in \mathcal{F}$, so $E_1 \in \mathcal{F}$. Hence E is $\mathcal{F}$ -subnormal in E_1 by Lemma 2.2. By hypothesis, E_1 is $\mathcal{F}$ -subnormal in G . Therefore E is $\mathcal{F}$ -subnormal in G . The lemma is proved.

Lemma 2.4 (See [16, Chapter VI, Theorem 24.2]). *Let $\mathcal{F}$ be a saturated formation and G a soluble group. If $G^{\mathcal{F}} \neq 1$ and every $\mathcal{F}$ -abnormal maximal subgroup of G belongs to $\mathcal{F}$, then the following hold:*

- (1) $G^{\mathcal{F}}$ is a p -group for some prime p ;
- (2) $G^{\mathcal{F}}/\Phi(G^{\mathcal{F}})$ is an $\mathcal{F}$ -eccentric chief factor of G ;
- (3) if $G^{\mathcal{F}}$ is a non-abelian group, then the center, commutator subgroup, and Frattini subgroup of G coincide and are of exponent p ;

- (4) if $G^{\mathcal{F}}$ is abelian, then $G^{\mathcal{F}}$ is elementary;
- (5) if $p > 2$, then $G^{\mathcal{F}}$ is of exponent p ; for $p = 2$ the exponent of $G^{\mathcal{F}}$ is at most 4;
- (6) every pair of $\mathcal{F}$ -abnormal maximal subgroups of G are conjugate in G .

Lemma 2.5 (See [16, Chapter VI, Theorem 24.5]). *Let $\mathcal{F}$ be a saturated formation. Let G be an $\mathcal{F}$ -critical group and G has a normal Sylow p -subgroup $G_p \neq 1$ for some prime p . Then:*

- (1) $G_p = G^{\mathcal{F}}$;
- (2) $F(G) = G_p \Phi(G)$;
- (3) $G_{p'} \cap C_G(G_p / \Phi(G_p)) = \Phi(G) \cap G_{p'}$, where $G_{p'}$ is some complement of G_p in G .

Lemma 2.6 (See [16, Chapter VI, Theorems 26.3 and 26.5]). *Let G be an $\mathcal{U}$ -critical group. Then:*

- (1) G is soluble and $|\pi(G)| \leq 3$;
- (2) if G is not a Schmidt group, then G is Ore dispersive;
- (3) $G^{\mathcal{U}}$ is the unique normal Sylow subgroup of G ;
- (4) if S is a complement of $G^{\mathcal{U}}$ in G , then $S/S \cap \Phi(G)$ is either a primary cyclic group or a Miller-Moreno group.

Recall that the product of all normal subgroups of a group G whose G -chief factors are $\mathcal{F}$ -central in G is called $\mathcal{F}$ -hypercentre of G and denoted by $Z_{\mathcal{F}}(G)$ [1, p. 389].

Lemma 2.7 (See [17, Lemma 2.14]). *Let $\mathcal{F}$ be a saturated formation and F the canonical local satellite of $\mathcal{F}$. Let E be a normal p -subgroup of a group G . Then $E \leq Z_{\mathcal{F}}(G)$ if and only if $G/C_G(E) \in F(p)$.*

The product $\mathcal{MH}$ of the formations $\mathcal{M}$ and $\mathcal{H}$ is the class of all groups G such that $G^{\mathcal{H}} \in \mathcal{M}$.

Lemma 2.8 (See [3, Corollary 7.14]). *The product of any two n -multiply saturated formations is an n -multiply saturated formation.*

We shall also need the following evident lemma.

Lemma 2.9. *If $G = AB$, then $G = AB^x$ for all $x \in G$.*

Let $\mathcal{F}$ be a class of groups and t a natural number with $t \geq 2$. Recall that $\mathcal{F}$ is called Σ_t -closed if $\mathcal{F}$ contains all such groups G that G has subgroups $H_1, \dots, H_t$ whose indices are pairwise coprime and $H_i \in \mathcal{F}$, for $i = 1, \dots, t$.

Lemma 2.10 (See [16, Chapter I, Lemma 4.11]). *Every formation of nilpotent groups is Σ_3 -closed.*

If $\mathcal{F} = LF(f)$ and $f(p) \subseteq \mathcal{F}$ for all primes p , then f is called an *integrated local satellite* of $\mathcal{F}$. Let $\mathcal{X}$ be a set of groups. The symbol $l_n \text{form} \mathcal{X}$ denotes the intersection of all n -multiply saturated formations $\mathcal{F}$ such that $\mathcal{X} \subseteq \mathcal{F}$. In view of [15, Remak 3.1.7], $l_n \text{form} \mathcal{X}$ is an n -multiply saturated

formation.

Lemma 2.11 (See [3, Theorem 8.3]). *Let $\mathcal{F}$ be an n -multiply saturated formation. Then $\mathcal{F}$ has an integrated local satellite f such that $f(p) = l_{n-1}\text{form}(G/O_{p',p}(G)|G \in \mathcal{F})$ for all primes p .*

Lemma 2.12 (See [18, Section 1.4]). *Every r -multiply saturated formation contained in $\mathcal{N}^{r+1}$ is hereditary.*

Lemma 2.13 (See [16, p. 35]). *For any ordering ϕ of $\mathbb{P}$ the class of all ϕ -dispersive groups is a saturated formation.*

Lemma 2.14 (See [17, Corollary 1.6]). *Let $\mathcal{F}$ be a saturated formation containing all nilpotent groups and E a normal subgroup of G . If $E/E \cap \Phi(G) \in \mathcal{F}$, then $E \in \mathcal{F}$.*

Lemma 2.15 (See [16, Theorem 15.10]). *Let $\mathcal{F}$ be a saturated formation and G a group such that $G^{\mathcal{F}}$ is nilpotent. Let H and M be subgroups of G , $H \in \mathcal{F}$, $H \leq M$ and $HF(G) = G$. If H is $\mathcal{F}$ -subnormal in M , then $M \in \mathcal{F}$.*

3 Proof of Theorem A

First we give two propositions which may be independently interesting since they generalize some known results.

Proposition 3.1. *Suppose that $G = A_1A_2 = A_2A_3 = A_1A_3$, where A_1 , A_2 and A_3 are soluble subgroups of G . If the indices $|G : N_G(A'_1)|$, $|G : N_G(A'_2)|$, $|G : N_G(A'_3)|$ are pairwise coprime, then G is soluble.*

Corollary 3.2. *Suppose that $G = A_1A_2 = A_2A_3 = A_1A_3$, where A_1 , A_2 and A_3 are soluble subgroups of G . If the indices $|G : N_G(A_1)|$, $|G : N_G(A_2)|$, $|G : N_G(A_3)|$ are pairwise coprime, then G is soluble.*

Corollary 3.3 (H. Wielandt). *If G has three soluble subgroups A_1 , A_2 and A_3 whose indices $|G : A_1|$, $|G : A_2|$, $|G : A_3|$ are pairwise coprime, then G is itself soluble.*

Proposition 3.4. *Let $\mathcal{M}$ be an r -multiply saturated formation and $\mathcal{N} \subseteq \mathcal{M} \subseteq \mathcal{N}^{r+1}$ for some $r \geq 0$. Then, for any prime p , both formations $\mathcal{M}$ and $\mathcal{G}_p\mathcal{M}$ are Σ_{r+3} -closed.*

Proof. Let M be the canonical local satellite of $\mathcal{M}$. Let $\mathcal{F}$ be one of the formations $\mathcal{M}$ or $\mathcal{G}_p\mathcal{M}$. Let G be any group such that for some subgroups $H_1, \dots, H_{r+3}$ of G whose indices $|G : H_1|, \dots, |G : H_{r+3}|$ are pairwise coprime we have $H_1, \dots, H_{r+3} \in \mathcal{F}$. We shall prove $G \in \mathcal{F}$. Suppose that this is false and let G be a counterexample with $r + |G|$ minimal. Let N be a minimal normal subgroup of G .

(1) $N = G^{\mathcal{F}}$ is the only minimal normal subgroup of G and $N \leq O_q(G)$ for some prime q . Hence if $\mathcal{F} = \mathcal{G}_p\mathcal{M}$, then $q \neq p$.

It is clear that the hypothesis holds for G/N , so $G/N \in \mathcal{F}$ by the choice of G . Hence $N = G^{\mathcal{F}}$ since $G \notin \mathcal{F}$. Moreover, N is a q -group for some prime q since G is soluble by Proposition 3.1.

Finally, if $\mathcal{F} = \mathfrak{G}_p\mathcal{M}$ and $p = q$, then

$$G \in \mathfrak{G}_p(\mathfrak{G}_p\mathcal{M}) = \mathfrak{G}_p\mathcal{F} = \mathcal{F},$$

a contradiction. Hence we have (1).

Since the indices $|G : H_1|, \dots, |G : H_{r+3}|$ are pairwise coprime, in view of (1) we may assume without loss of generality that $N \leq H_i$ for all $i = 2, \dots, r+3$.

$$(2) \ C_G(N) = N.$$

First we show that $N \not\leq \Phi(G)$. Suppose that $N \leq \Phi(G)$. If $r > 0$, then $\mathcal{F}$ is saturated by Lemma 2.8, so $G \in \mathcal{F}$. This contradiction shows that $r = 0$ and so $\mathcal{F} = \mathfrak{G}_p\mathcal{M}$ by Lemma 2.10 and the choice of G . Hence $q \neq p$ by (1). Let $O/N = O_p(G/N)$ and P be a Sylow p -subgroup of O . Then $G = ON_G(P) = NPN_G(P) = NN_G(P) = N_G(P)$ by the Frattini Argument since $N \leq \Phi(G)$. Hence in view of (1), $O_p(G/N) = 1$ and so $G/N \in \mathcal{M}$ since $G/N \in \mathcal{F} = \mathfrak{G}_p\mathcal{M}$. But then G is a p' -group. Hence $H_1, H_2, H_3 \in \mathcal{M}$. Thus $G \in \mathcal{M} \subseteq \mathcal{F}$ by Lemma 2.10. This contradiction shows that $N \not\leq \Phi(G)$. But then $C_G(N) = N$ by (1) and [1, A, Theorem 15.2].

$$(3) \ r > 0.$$

Suppose that $r = 0$. Then $\mathcal{F} = \mathfrak{G}_p\mathcal{M}$, where $\mathcal{M}$ is a formation of nilpotent groups. Since $N \leq H_2 \in \mathcal{F}$ and, by (2), $C_G(N) = N$, $O_p(H_2) = 1$. Hence H_2 is a p' -group. Similarly, H_3 is a p' -group. Hence $G = H_1H_2$ is a p' -group. But then $H_1 \in \mathcal{M}$, so $G \in \mathcal{F}$ by Lemma 2.10. This contradiction shows that we have (3).

$$(4) \ H_i/N \in M(q) \text{ for all } i = 2, \dots, r+3.$$

Let $i \in \{2, \dots, r+3\}$. Then $H_i \in \mathcal{M}$. Indeed, if $\mathcal{F} = \mathfrak{G}_p\mathcal{M}$, then $q \neq p$ by (1). On the other hand, in view of (2), $C_G(N) = N$. Hence $O_p(H_i) = 1$, which implies that $H_i \in \mathcal{M}$. But then, by (2) and Lemma 2.7, $H_i/N = H_i/C_{H_i}(N) \in M(q)$.

$$(5) \ G/N \in M(q).$$

By Lemma 2.11 and [1, Chapter IV, Proposition 3.8], $M(q) = \mathfrak{G}_q\mathcal{M}_0$, where $\mathcal{M}_0 = l_{r-1}\text{form}$ ($G/O_{q',q}(G) \mid G \in \mathcal{M}$). Since $\mathcal{M} \subseteq \mathcal{N}^{r+1}$, $G/O_{q',q}(G) \in \mathcal{N}^r$, so $\mathcal{M}_0 \subseteq \mathcal{N}^r$ since $\mathcal{M}_0$ is an $(r-1)$ -multiply saturated formation. Therefore, the minimality of $r + |G|$ and Claim (4) imply that $G/N \in M(q)$.

Final contradiction. Since N is a q -group by (1), from (5) it follows that $G \in \mathfrak{G}_qM(q) = M(q) \subseteq \mathcal{M} \subseteq \mathfrak{G}_p\mathcal{M}$. This contradiction completes the proof of the proposition.

Corollary 3.5 (See [20, Satz 1.3]). *Every saturated formation contained in $\mathcal{N}^2$ is Σ_4 -closed.*

Corollary 3.6. *The class of all soluble groups of nilpotent length at most r ($r \geq 2$) is Σ_{r+2} -closed.*

Proof. It is clear that $\mathcal{N}^r$ is hereditary formation. Moreover, in view of Lemma 2.8, $\mathcal{N}^r$ is an $(r-1)$ -multiply saturated formation. So $\mathcal{N}^r$ is Σ_{r+2} -closed by Proposition 3.4.

Proof of Theorem A. Suppose that the theorem is false and consider some counterexample

G of minimal order. Take a maximal subgroup M of G . Then by hypothesis all $(n-1)$ -maximal subgroups of M are $\mathcal{F}$ -subnormal in G , and so they are $\mathcal{F}$ -subnormal in M by Lemmas 2.1(1) and 2.12. The solubility of G implies that either $|\pi(M)| = |\pi(G)|$ or $|\pi(M)| = |\pi(G)| - 1$, so $M \in \mathcal{F}$ by the choice of G . Hence G is an $\mathcal{F}$ -critical group.

Since G is soluble, G has a maximal subgroup T with $|G : T| = p^a$ for any prime p dividing $|G|$. On the other hand, $\mathcal{F}$ is Σ_{r+3} -closed by Proposition 3.4. Hence $|\pi(G)| \leq r+2$. Moreover, by hypothesis, $|\pi(G)| \geq n+r+1$. Therefore $n=1$. Thus all maximal subgroups of G are $\mathcal{F}$ -normal, so $G/\Phi(G) \in \mathcal{F}$. But $\mathcal{F}$ is a saturated formation and hence $G \in \mathcal{F}$. This contradiction completes the proof of the result.

Corollary 3.7 (See [14, Theorem 6]). *If each n -maximal subgroup of a soluble group G is subnormal, and if $|\pi(G)| \geq n+1$, then G is nilpotent.*

Corollary 3.8 (See [21, Theorem A]). *If every n -maximal subgroup of a soluble group G is $\mathcal{U}$ -subnormal in G and $|\pi(G)| \geq n+2$, then G is supersoluble.*

Corollary 3.9. *Let $\mathcal{F}$ be the class of all groups G with $G' \leq F(G)$. If every n -maximal subgroup of a soluble group G is $\mathcal{F}$ -subnormal in G and $|\pi(G)| \geq n+2$, then $G \in \mathcal{F}$.*

Corollary 3.10. *If every n -maximal subgroup of a soluble group G is $\mathcal{N}^r$ -subnormal in G ($r \geq 1$) and $|\pi(G)| \geq n+r$, then $G \in \mathcal{N}^r$.*

4 Proofs of Theorems B, C, and D

Proof of Theorem C. First suppose that every 2-maximal subgroup of G is $\mathcal{F}$ -subnormal in G . Assume that $G \notin \mathcal{F}$. We shall show that G is an $\mathcal{F}$ -critical group and $G^{\mathcal{F}}$ is a minimal normal subgroup of G . Let M be a maximal subgroup of G and T a maximal subgroup of M . By hypothesis, T is $\mathcal{F}$ -subnormal in G . Therefore T is $\mathcal{F}$ -normal in M by Lemma 2.1(1), so $M/T_M \in \mathcal{F}$. Since T is arbitrary and $\mathcal{F}$ is saturated, $M \in \mathcal{F}$. Consequently, all maximal subgroups of G belong to $\mathcal{F}$. Hence G is an $\mathcal{F}$ -critical group. Then by hypothesis, G is soluble and it has a normal Sylow p -subgroup $G_p \neq 1$ for some prime p . Thus $G_p = G^{\mathcal{F}}$ by Lemma 2.5. On the other hand, by Lemma 2.4, $G_p/\Phi(G_p)$ is a chief factor of G .

Let M be an $\mathcal{F}$ -abnormal maximal subgroup of G . Then $G_p \not\leq M$, so $G = G_p M$ and $M = (G_p \cap M)G_{p'} = \Phi(G_p)G_{p'}$, where $G_{p'}$ is a Hall p' -subgroup of G . Assume that $\Phi(G_p) \neq 1$. It is clear that $\Phi(G_p) \not\leq \Phi(M)$. Let T be a maximal subgroup of M such that $\Phi(G_p) \not\leq T$. Then $M = \Phi(G_p)T$. Since T is $\mathcal{F}$ -subnormal in G , there is a maximal subgroup L of G such that $T \leq L$ and $G/L_G \in \mathcal{F}$. Then $G_p \leq L_G$, so $G = G_p M = G_p \Phi(G_p)T = G_p T \leq L$, a contradiction. Hence $\Phi(G_p) = 1$. Therefore $G_p = G^{\mathcal{F}}$ is a minimal normal subgroup of G by Lemma 2.4.

Now suppose that G is an $\mathcal{F}$ -critical group and $G^{\mathcal{F}}$ is a minimal normal subgroup of G . Let T be a 2-maximal subgroup of G and M a maximal subgroup of G such that T is a maximal subgroup

of M . Since $M \in \mathcal{F}$, T is $\mathcal{F}$ -subnormal in M by Lemma 2.2. Therefore, if M is $\mathcal{F}$ -normal in G , then T is $\mathcal{F}$ -subnormal in G by Lemma 2.1(3). Assume that M is $\mathcal{F}$ -abnormal in G . Then $G^{\mathcal{F}} \not\leq M$. Therefore, since $G^{\mathcal{F}}$ is a minimal normal subgroup of G by hypothesis, $G = G^{\mathcal{F}} \rtimes M$ and $G^{\mathcal{F}}T$ is a maximal $\mathcal{F}$ -normal subgroup of G . Moreover, since G is an $\mathcal{F}$ -critical group, $G^{\mathcal{F}}T \in \mathcal{F}$ and hence T is $\mathcal{F}$ -subnormal in $G^{\mathcal{F}}T$ by Lemma 2.2. Hence, T is $\mathcal{F}$ -subnormal in G . The theorem is proved.

From Theorem C and Lemma 2.6 we get

Corollary 4.1 (See [21, Theorem 3.1]). *Every 2-maximal subgroup of G is $\mathcal{U}$ -subnormal in G if and only if G is an $\mathcal{U}$ -critical group and $G^{\mathcal{U}}$ is a minimal normal subgroup of G .*

Proof of Theorem B. First suppose that all n -maximal subgroups of G are $\mathcal{F}$ -subnormal in G . We shall show, in this case, that either $G \in \mathcal{F}$ or G is a group of the type II. Assume that this is false and consider a counterexample G for which $|G| + n$ is minimal. Therefore $A = G^{\mathcal{F}} \neq 1$. Then:

(a) *The hypothesis holds for every maximal subgroup of G .*

Let M be a maximal subgroup of G . Then by hypothesis, all $(n-1)$ -maximal subgroups of M are $\mathcal{F}$ -subnormal in G , and so they are $\mathcal{F}$ -subnormal in M by Lemmas 2.1(1) and 2.12. Moreover, the solubility of G implies that either $|\pi(M)| = |\pi(G)|$ or $|\pi(M)| = |\pi(G)| - 1$.

(b) *If M is a maximal subgroup of G and $|\pi(M)| = |\pi(G)|$, then $M \in \mathcal{F}$.*

In view of hypothesis and Lemmas 2.1(1) and 2.12, all $(n-1)$ -maximal subgroups of M are $\mathcal{F}$ -subnormal in M . Since $|\pi(M)| = |\pi(G)| \geq n+1 = n-1+2$, $M \in \mathcal{F}$ by Theorem A.

(c) *If W is a Hall q' -subgroup of G for some $q \in \pi(G)$, then either $W \in \mathcal{F}$ or W is a group of the type II.*

If W is not a maximal subgroup of G , then there is a maximal subgroup V of G such that $W \leq V$ and $|\pi(V)| = |\pi(G)|$. By (b), $V \in \mathcal{F}$. Hence $W \in \mathcal{F}$ by Lemma 2.12. Suppose that W is a maximal subgroup of G . Then by (a), the hypothesis holds for W , so either $W \in \mathcal{F}$ or W is a group of the type II by the choice of G .

(d) *The hypothesis holds for G/N , where N is a minimal normal subgroup of G .*

If N is not a Sylow subgroup of G , then $|\pi(G/N)| = |\pi(G)|$. Moreover, if H/N is an n -maximal subgroup of G/N , then H is an n -maximal subgroup of G . Therefore H is $\mathcal{F}$ -subnormal in G . Consequently, H/N is $\mathcal{F}$ -subnormal in G/N by Lemma 2.1(2). But if G/N has no n -maximal subgroups, then by the solubility of G , the identity subgroup of G/N is $\mathcal{F}$ -subnormal in G/N and it is the unique i -maximal subgroup of G/N for some $i < n$ with $i < |\pi(G/N)|$. Finally, consider the case that N is a Sylow p -subgroup of G . Let E be a Hall p' -subgroup of G . It is clear that $|\pi(E)| = |\pi(G)| - 1$ and E is a maximal subgroup of G .

Let H/N be an $(n-1)$ -maximal subgroup of G/N . Then H is an $(n-1)$ -maximal subgroup of G and $H = H \cap NE = N(H \cap E)$. There is a chain of subgroups $H = H_0 < H_1 < \dots < H_{n-1} = G$ of G , where H_{i-1} is a maximal subgroup of H_i ($i = 1, \dots, n-1$). Then $H_{i-1} \cap E$ is a maximal

subgroup of $H_i \cap E$, for $i = 1, \dots, n-1$. Indeed, suppose that for some i there is a subgroup K of $H_i \cap E$ such that $H_{i-1} \cap E \leq K \leq H_i \cap E$. Then $(H_{i-1} \cap E)N \leq KN \leq (H_i \cap E)N$, so $H_{i-1} = H_{i-1} \cap EN \leq KN \leq H_i \cap EN = H_i$. Whence either $KN = H_{i-1}$ or $KN = H_i$. If $KN = H_{i-1}$, then $H_{i-1} \cap E = KN \cap E = K(N \cap E) = K$. In the second case we have $H_i \cap E = KN \cap E = K(N \cap E) = K$. Therefore $H_{i-1} \cap E$ is a maximal subgroup of $H_i \cap E$, so $H \cap E$ is an $(n-1)$ -maximal subgroup of E . Since E is a maximal subgroup of G , $H \cap E$ is an n -maximal subgroup of G . Hence $H \cap E$ is $\mathcal{F}$ -subnormal in G by hypothesis. Therefore $H/N = (H \cap E)N/N$ is $\mathcal{F}$ -subnormal in G/N by Lemma 2.1(2).

(e) $|\pi(G)| > 2$.

If $|\pi(G)| = 2$, then $n = 1$ and so all maximal subgroups of G are $\mathcal{F}$ -normal by hypothesis. Hence $G \in \mathcal{F}$ since $\mathcal{F}$ is a saturated formation, a contradiction.

(f) G is an Ore dispersive group.

Suppose that this is false. Take a minimal normal subgroup N of G . Then by (d), the hypothesis holds for G/N , so either $G/N \in \mathcal{F}$ or G/N is a group of the type II. Thus, in view of $\mathcal{F} \subseteq \mathcal{U}$ and the choice of G , G/N is an Ore dispersive group. By Lemma 2.13, the class of all Ore dispersive groups is a saturated formation. Therefore N is the unique minimal normal subgroup of G and $N \not\leq \Phi(G)$. Hence $\Phi(G) = 1$ and there is a maximal subgroup L of G such that $G = N \rtimes L$ and $L_G = 1$. Thus $C_G(N) = N$ by [1, A, Theorem 15.2].

Since G is soluble, G has a normal maximal subgroup M with $|G : M| = p$ for some prime p and either $|\pi(M)| = |\pi(G)|$ or $|\pi(M)| = |\pi(G)| - 1$. By (a), the hypothesis holds for M . Therefore, in view of $\mathcal{F} \subseteq \mathcal{U}$ and the choice of G , M is an Ore dispersive group. Denote by q the greatest number in $\pi(M)$. Take a Sylow q -subgroup M_q of M . Since M_q is a characteristic subgroup of M , M_q is normal in G . Consider the case $|\pi(M)| = |\pi(G)|$ first. Then q is the greatest prime divisor of the order of G and $M_q \neq 1$. Hence G/M_q is an Ore dispersive group, and by the maximality of q , so is G . Suppose now that $|\pi(M)| = |\pi(G)| - 1$. If $q > p$, then, as above, we conclude that G is an Ore dispersive group as well. Let $p > q$. Then p is the greatest prime divisor of $|G|$. Since $M_q \neq 1$, it follows that $N \leq M_q$, so N is a q -group. In addition, since $|\pi(G)| > 2$ by (e), there is a prime divisor r of the order of G such that $q \neq r \neq p$. Take a Hall r' -subgroup W of G . Then $PN \leq W$ for some Sylow p -subgroup P of G . Moreover, by (c), W is an Ore dispersive group. Hence P is normal in W , and so $P \leq C_G(N) = N$. The resulting contradiction shows that G is an Ore dispersive group.

(g) A is a nilpotent group.

Suppose that this is false. Let N be a minimal normal subgroup of G . Then by (d), $(G/N)^{\mathcal{F}} = G^{\mathcal{F}}N/N \simeq G^{\mathcal{F}}/G^{\mathcal{F}} \cap N$ is a nilpotent group. It is known that the class of all nilpotent groups is a saturated formation. Hence in the case when G has a minimal normal subgroup $R \neq N$ we have $G^{\mathcal{F}}/(G^{\mathcal{F}} \cap N) \cap (G^{\mathcal{F}} \cap R) \simeq G^{\mathcal{F}}$ is nilpotent. Thus N is the unique minimal normal subgroup of G and $N \leq G^{\mathcal{F}}$. If $N \leq \Phi(G)$, then $G^{\mathcal{F}}/G^{\mathcal{F}} \cap \Phi(G) \simeq (G^{\mathcal{F}}/N)/((G^{\mathcal{F}} \cap \Phi(G))/N)$ is nilpotent, so $G^{\mathcal{F}}$ is nilpotent by Lemma 2.14. Therefore $N \not\leq \Phi(G)$. Hence $\Phi(G) = 1$ and there is a maximal subgroup

L of G such that $G = N \rtimes L$ and $L_G = 1$. Thus $C_G(N) = N$ by [1, A, Theorem 15.2] and $N \neq A$.

Case 1: $|\pi(G)| = 3$. By hypothesis, either all maximal subgroups of G or all its 2-maximal subgroups are $\mathcal{F}$ -subnormal in G . In the first case we infer that $G \in \mathcal{F}$, which contradicts the choice of G . Hence all 2-maximal subgroups of G are $\mathcal{F}$ -subnormal. Since $\mathcal{F} \subseteq \mathcal{U}$, in view of Lemma 2.6, every $\mathcal{F}$ -critical group has a normal Sylow subgroup. Whence Theorem C implies that G is an $\mathcal{F}$ -critical group and $A = G^{\mathcal{F}}$ is a minimal normal subgroup of G . Therefore $A = N$, a contradiction.

Case 2: $|\pi(G)| \geq 4$. Assume that N is a p -group, and take a Sylow subgroup P of G such that $N \leq P$. Observe that if $N \neq P$, then $L \in \mathcal{F}$ by (b), and so $A = N$, a contradiction. Hence $N = P$.

Case 2.1: $|\pi(G)| = 4$.

(1) *All 3-maximal subgroups of G are $\mathcal{F}$ -subnormal in G and L is an $\mathcal{F}$ -critical group.*

Since $G \notin \mathcal{F}$ and $|\pi(G)| = 4$, either all 2-maximal subgroups of G or all its 3-maximal subgroups are $\mathcal{F}$ -subnormal in G . In the first case G is an $\mathcal{F}$ -critical group and $A = G^{\mathcal{F}}$ is a minimal normal subgroup of G by Theorem C. Hence $A = N$, a contradiction. Therefore all 3-maximal subgroups of G are $\mathcal{F}$ -subnormal in G . Thus all second maximal subgroups of G belong to $\mathcal{F}$ by Lemma 2.3. Consequently, either $L \in \mathcal{F}$ or L is an $\mathcal{F}$ -critical group. But in the first case $N = A$, a contradiction. Therefore L is an $\mathcal{F}$ -critical group.

(2) *$L = Q \rtimes (R \rtimes T)$, where Q, R, T are Sylow subgroups of G , $Q = L^{\mathcal{F}}$ is a minimal normal subgroup of L , and $G^{\mathcal{F}} = PQ$.*

Since $N = P$ is a Sylow p -subgroup of G and $|\pi(G)| = 4$, $|\pi(L)| = 3$. Hence in view of (f), $L = Q \rtimes (R \rtimes T)$, where Q, R, T are Sylow subgroups of G . Moreover, $Q = L^{\mathcal{F}}$ by Lemma 2.5 and Q is a minimal normal subgroup of L by Theorem C since every 2-maximal subgroup of L is $\mathcal{F}$ -subnormal in L by (1) and Lemmas 2.1(1) and 2.12. Finally, since $G/N \notin \mathcal{F}$ and $G/PQ \simeq L/Q \in \mathcal{F}$, we have $G^{\mathcal{F}} = PQ$.

(3) *$V = PQR$ is not supersoluble. Hence $V \notin \mathcal{F}$.*

Assume that V is a supersoluble group. Since $F(V)$ is a characteristic subgroup of V and V is a normal subgroup of G , $F(V)$ is normal in G . Hence every Sylow subgroup of $F(V)$ is normal in G . But N is the unique minimal normal subgroup of G . Therefore $F(V) = N = P$. Thus $V/P \simeq QR$ is an abelian group. Hence R is normal in L and so $R \leq F(L)$. In view of Lemma 2.5, $F(L) = Q\Phi(L)$. Whence $R \leq \Phi(L)$. This contradiction shows that V is not supersoluble. Thus $V \notin \mathcal{F}$ since $\mathcal{F} \subseteq \mathcal{U}$ by hypothesis.

(4) *V is a maximal subgroup of G . Hence $|T| = t$ is a prime.*

If V is not a maximal subgroup of G , then there is a maximal subgroup U of G such that $V \leq U$ and $|\pi(U)| = |\pi(G)|$. Hence $U \in \mathcal{F}$ by (b), so $V \in \mathcal{F}$ by Lemma 2.12, a contradiction. Therefore V is a normal maximal subgroup of G . Whence $|T|$ is a prime.

(5) *$|Q| = q$ is a prime and $R = \langle x \rangle$ is a cyclic group.*

Since V is a maximal subgroup of G by (4), all 2-maximal subgroups of V are $\mathcal{F}$ -subnormal in V by (1) and Lemmas 2.1(1) and 2.12. Hence, in view of (3), V is an $\mathcal{F}$ -critical group by Theorem C. Therefore, in fact, V is an $\mathcal{U}$ -critical group by (3) since $\mathcal{F} \subseteq \mathcal{U}$. Hence QR is supersoluble. Since V is normal in G and $\Phi(G) = 1$, $\Phi(V) = 1$. Therefore QR is a Schmidt group by Lemma 2.6. Hence R is cyclic and Q is a minimal normal subgroup of QR by Lemma 2.4. Whence $|Q|$ is a prime.

(6) $|R| = r$ is a prime and $C_L(Q) = Q$.

By (4) and (5), L is a supersoluble group. Suppose that $|R| = r^b$ is not a prime and let M be a maximal subgroup of L such that $|L : M| = r$. Let $W = PM$. Then $\pi(W) = \pi(G)$, so $W \in \mathcal{F}$ by (b) and hence W is supersoluble. Since $C_G(N) = N$, $F(W) = P$. Hence $W/P \simeq M$ is abelian. It is clear that $Q \leq M$, so $M \leq C_L(Q)$. Hence $T \leq F(L)$. On the other hand, $F(L) = Q\Phi(L)$ by Lemma 2.5. Therefore $T \not\leq F(L)$. This contradiction shows that $|R| = r$ and so $C_L(Q) = Q$ by Lemma 2.5.

(7) $1 \neq C_G(x) \cap PQ = P_1 \leq P$.

Suppose that $C_G(x) \cap PQ = 1$. Then by the Thompson's theorem [22, Theorem 10.5.4], PQ is a nilpotent group, so $Q \leq C_G(P) = P$, a contradiction. Thus $C_G(x) \cap PQ \neq 1$. Suppose that q divides $|C_G(x) \cap PQ|$. Then, by (5), for some $a \in P$ we have $Q^a \leq C_G(x) \cap PQ$, so $\langle Q^a, RT \rangle \leq N_G(R)$. Hence if E is a Hall p' -subgroup of $N_G(R)$, then $E \simeq L$. Therefore L has a normal r -subgroup, so $C_L(Q) \neq Q$, a contradiction. Thus $C_G(x) \cap PQ = P_1 \leq P$.

Final contradiction for Case 2.1. Let $D = \langle P_1, RT \rangle$. Then $D \leq N_G(R)$. If q divides $|D|$, then, as above, we have $C_L(Q) \neq Q$. Thus $D \cap Q^a = 1$ for all $a \in P$. Moreover, if $P \leq D$, then $PR = P \times R$ and $R \leq C_G(P) = P$. Therefore $P \not\leq D$ and D is not a maximal subgroup of G . Hence D is a k -maximal subgroup of G for some $k \geq 2$. Then there is a 3-maximal subgroup S of G such that $RT \leq S \leq D$. By hypothesis, S is $\mathcal{F}$ -subnormal in G . Hence at least one of the maximal subgroups L or PRT is $\mathcal{F}$ -normal in G , contrary to (2).

Case 2.2: $|\pi(G)| > 4$. If $\pi(L) = \{p_1, \dots, p_t\}$, then $t > 3$. Let E_i be a Hall p'_i -subgroup of L and $X_i = PE_i$. We shall show that $E_i \in \mathcal{F}$ for all $i = 1, \dots, t$. By (c), either $X_i \in \mathcal{F}$ or X_i is a group of the type II, for $i = 1, \dots, t$. In the former case we have $E_i \simeq X_i/P \in \mathcal{F}$. Assume that X_i be a group of the type II. Then $X_i^{\mathcal{F}}$ is nilpotent, so $X_i^{\mathcal{F}} \leq F(X_i)$. But since P is normal in X_i and $C_G(P) = P$, $F(X_i) = P$. Hence $X_i^{\mathcal{F}} = P$, so $E_i \in \mathcal{F}$. Since $t > 3$, Proposition 3.4 implies that then $L \in \mathcal{F}$. Therefore $A = N$, a contradiction. Hence we have (g).

(h) A is a Hall subgroup of G .

Suppose that this is false. Since G is Ore dispersive by (f), for the greatest prime divisor p of $|G|$ the Sylow p -subgroup P is normal in G . Assume that P is not a minimal normal subgroup of G . Then there is a maximal subgroup M of G such that $G = PM$ and $P \cap M \neq 1$. Since $|\pi(M)| = |\pi(G)|$, $M \in \mathcal{F}$ by (b). Hence $G/P \simeq M/M \cap P \in \mathcal{F}$, so $A = G^{\mathcal{F}} \leq P$. Suppose that $\Phi(P) \neq 1$. Let N be a minimal normal subgroup of G such that $N \leq \Phi(P)$. By (d), the hypothesis holds for G/N , so either $G/N \in \mathcal{F}$ or G/N is a group of the type II by the choice of G . If $G/N \in \mathcal{F}$,

then $A = N \leq \Phi(P)$. Since P is normal in G , $\Phi(P) \leq \Phi(G)$. Thus $A \leq \Phi(G)$ and so $G \in \mathcal{F}$, a contradiction. Hence G/N is a group of the type II. Therefore $AN/N = G^{\mathcal{F}}N/N = (G/N)^{\mathcal{F}}$ is a Hall subgroup of G/N . Consequently, $AN = P$. Hence $A\Phi(P) = P$, so $A = P$, a contradiction. Thus $\Phi(P) = 1$. By Maschke's theorem, $P = N_1 \times \dots \times N_k$ is the direct product of some minimal normal subgroups of G . If $N_1 \neq P$, then $G/N_1 \in \mathcal{F}$ and $G/N_2 \in \mathcal{F}$ by Theorem A. Consequently, so is G . This contradiction shows that P is a minimal normal subgroup of G .

By (d), the hypothesis holds for G/P , so either $G/P \in \mathcal{F}$ or G/P is a group of the type II by the choice of G . If $G/P \in \mathcal{F}$, then $A = P$, a contradiction. Hence G/P is a group of the type II. Therefore $AP/P = G^{\mathcal{F}}P/P = (G/P)^{\mathcal{F}}$ is a Hall subgroup of G/P . If $P \leq A$, then $A = P \rtimes A_{p'}$, where $A_{p'}$ is a Hall p' -subgroup of A . But since $A_{p'} \simeq A/P$ and A/P is a Hall subgroup of G/P , A is a Hall subgroup of G . Therefore $P \cap A = 1$, so A is a Hall subgroup of G since $AP/P \simeq A/A \cap P \simeq A$.

(i) A is either of the form $N_1 \times \dots \times N_t$, where each N_i is a minimal normal subgroup of G , which is a Sylow subgroup of G , for $i = 1, \dots, t$, or a Sylow p -subgroup of G of exponent p for some prime p and the commutator subgroup, the Frattini subgroup, and the center of A coincide, while $A/\Phi(A)$ is an $\mathcal{F}$ -eccentric chief factor of G .

Suppose that A is not a minimal normal subgroup of G . Take a Sylow p -subgroup P of A , where p divides $|A|$. Claims (g) and (h) imply that P is a normal Sylow subgroup of G . Let N be a minimal normal subgroup of G with $N \leq P$. First suppose that $N \leq \Phi(G)$, and take a maximal subgroup M of G with $P \not\leq M$. Then $M \in \mathcal{F}$ by (b). Therefore $G/P \simeq M/M \cap P \in \mathcal{F}$. In this case $A = P$. Moreover, if S is a maximal subgroup of G such that $P \not\leq S$, then $S \in \mathcal{F}$. Observe also that for every maximal subgroup X of G with $P \leq X$ we have X is $\mathcal{F}$ -subnormal in G . Thus, by Lemma 2.4, $A = G^{\mathcal{F}}$ satisfies condition II(1).

Suppose that for every minimal normal subgroup R of G such that $R \leq A$ we have $R \not\leq \Phi(G)$. Then there is a maximal subgroup L of G such that $G = N \rtimes L$. If $N \neq P$, then $L \in \mathcal{F}$ by (b). Therefore $A = N$, a contradiction. Consequently, all Sylow subgroups of A are minimal normal subgroups of G . Therefore $A = N_1 \times \dots \times N_t$, where N_i is a minimal normal subgroup of G , for $i = 1, \dots, t$.

(j) Every n -maximal subgroup of G belongs to $\mathcal{F}$ and induces on the Sylow p -subgroup of A the automorphism group which is contained in $F(p)$ for every prime divisor p of $|A|$.

Let H be any n -maximal subgroup of G . Suppose that H is a maximal subgroup of V , where V is an $(n-1)$ -maximal subgroup of G . Since $V \in \mathcal{F}$ by Lemmas 2.3 and 2.12, $H \in \mathcal{F}$.

Let $E = AH$. Since A is normal in E and A is nilpotent by (g), $A \leq F(E)$. Whence $E = F(E)H$. Since H is $\mathcal{F}$ -subnormal in G , H is $\mathcal{F}$ -subnormal in E by Lemmas 2.1(1) and 2.12. Moreover, $H \in \mathcal{F}$. Therefore $E \in \mathcal{F}$ by Lemma 2.15. Let P be a Sylow p -subgroup of A and K/L a chief factor of E such that $1 \leq L < K \leq P$. Since $E \in \mathcal{F}$, $E/C_E(K/L) \in F(p)$. Hence $P \leq Z_{\mathcal{F}}(E)$, so $E/C_E(P) \in F(p)$ by Lemma 2.7. Then $H/C_H(P) = H/C_E(P) \cap H \simeq HC_E(P)/C_E(P) \in F(p)$.

Now suppose that either $G \in \mathcal{F}$ or G is a group of type II. If $G \in \mathcal{F}$, then every subgroup of G is $\mathcal{F}$ -subnormal in G by Lemma 2.2. Let G be a group of type II. Take an n -maximal subgroup H of G . Put $E = G^{\mathcal{F}}H$. Let P be a Sylow p -subgroup of $G^{\mathcal{F}}$ and K/L a chief factor of E such that $1 \leq L < K \leq P$. By hypothesis, $H/C_H(P) \in F(p)$, so $H/C_H(K/L) \simeq (H/C_H(P))/(C_H(K/L)/C_H(P)) \in F(p)$. Since $G^{\mathcal{F}}$ is normal in E and $G^{\mathcal{F}}$ is nilpotent, $G^{\mathcal{F}} \leq F(E) \leq C_E(K/L)$. Hence

$$\begin{aligned} E/C_E(K/L) &= E/C_E(K/L) \cap E = E/C_E(K/L) \cap G^{\mathcal{F}}H = E/G^{\mathcal{F}}(C_E(K/L) \cap H) = \\ &= G^{\mathcal{F}}H/G^{\mathcal{F}}C_H(K/L) \simeq H/G^{\mathcal{F}}C_H(K/L) \cap H = H/C_H(K/L)(G^{\mathcal{F}} \cap H) = H/C_H(K/L), \end{aligned}$$

so $E/C_E(K/L) \in F(p)$ since $F(p)$ is hereditary by Lemma 2.12 and [15, Proposition 3.1.40]. Then $P \leq Z_{\mathcal{F}}(E)$, whence $G^{\mathcal{F}} \leq Z_{\mathcal{F}}(E)$. Thus $E/Z_{\mathcal{F}}(G) \in \mathcal{F}$. Hence $E \in \mathcal{F}$, so H is an $\mathcal{F}$ -subnormal subgroup of $G^{\mathcal{F}}H = E$. Since $G^{\mathcal{F}} \leq G^{\mathcal{F}}H$, $G^{\mathcal{F}}H$ is $\mathcal{F}$ -subnormal in G by Lemma 2.1(4). Consequently, in view of Lemma 2.1(3), H is $\mathcal{F}$ -subnormal in G . The theorem is proved.

Corollary 4.2 (See [21, Theorem B]). *Given a soluble group G with $|\pi(G)| \geq n+1$, all n -maximal subgroups of G are $\mathcal{U}$ -subnormal in G if and only if G is a group of one of the following types:*

I. G is supersoluble.

II. $G = A \rtimes B$, where $A = G^{\mathcal{U}}$ and B are Hall subgroups of G , while G is Ore dispersive and satisfies the following:

(1) A is either of the form $N_1 \times \dots \times N_t$, where each N_i is a minimal normal subgroup of G , which is a Sylow subgroup of G , for $i = 1, \dots, t$, or a Sylow p -subgroup of G of exponent p for some prime p and the commutator subgroup, the Frattini subgroup, and the center of A coincide, every chief factor of G below $\Phi(G)$ is cyclic, while $A/\Phi(A)$ is a noncyclic chief factor of G ;

(2) for every prime divisor p of the order of A every n -maximal subgroup H of G is supersoluble and induces on the Sylow p -subgroup of A an automorphism group which is an extension of some p -group by abelian group of exponent dividing $p-1$.

Proof of Theorem D. Assume that this is false and consider a counterexample G for which $|G| + n$ is minimal.

(a) G has a unique minimal normal subgroup N such that $C_G(N) = N$ and N is not a Sylow subgroup of G .

Let N be a minimal normal subgroup of G . Then the hypothesis holds for G/N (see Claim (d) in the proof of Theorem B). Consequently, G/N is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$ by the choice of G . Therefore N is not a Sylow subgroup of G . Moreover, by Lemma 2.13, $N \not\leq \Phi(G)$. Therefore G has a maximal subgroup M such that $G = N \rtimes M$. By Lemmas 2.1(1) and 2.12 all $(n-1)$ -maximal subgroups of M are $\mathcal{F}$ -subnormal in M . Moreover, $|\pi(M)| = |\pi(G)|$. Therefore Theorem B implies that $G/N \simeq M$ is an Ore dispersive group. Hence in the case when G has a minimal normal subgroup $R \neq N$ we have $G/N \cap R \simeq R$ is an Ore dispersive group. Thus N is the unique minimal normal subgroup of G , and so $C_G(N) = N$ by [1, A, Theorem 15.2].

(b) If W is a Hall q' -subgroup of G for some $q \in \pi(G)$, then W is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$.

If W is not a maximal subgroup of G , then there is a maximal subgroup V of G such that $W \leq V$ and $|\pi(W)| = |\pi(G)|$. By hypothesis, every $(n-1)$ -maximal subgroup of V is $\mathcal{F}$ -subnormal in G , so it is $\mathcal{F}$ -subnormal in V by Lemmas 2.1(1) and 2.12. Then, in view of Theorem B, V is Ore dispersive. Hence W is Ore dispersive. Suppose that W is a maximal subgroup of G . Then $|\pi(W)| = |\pi(G)| - 1$ and every $(n-1)$ -maximal subgroup of W is $\mathcal{F}$ -subnormal in W in view of hypothesis and Lemmas 2.1(1) and 2.12. Therefore W is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$ by the choice of G .

(c) $|\pi(G)| > 2$.

Suppose that $|\pi(G)| = 2$. Then by hypothesis, either all maximal subgroups of G or all its 2-maximal subgroups are $\mathcal{F}$ -subnormal in G . Therefore every maximal subgroup of G belongs to $\mathcal{F}$ in view of Lemmas 2.3 and 2.12. Consequently, either $G \in \mathcal{F}$ or G is an $\mathcal{F}$ -critical group. Since $\mathcal{F} \subseteq \mathcal{U}$, G is either a supersoluble group or an $\mathcal{U}$ -critical group. Therefore, in view of Lemma 2.6, G is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$, a contradiction.

Final contradiction. Suppose that N is a p -group, and take a prime divisor q of $|G|$ such that $q \neq p$. Take a Hall q' -subgroup E of G . Then $N \leq E$. By (b), E is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$. Consequently, some Sylow subgroup R of E is normal in E . Furthermore, if $N \not\leq R$, then $R \leq C_G(N) = N$. Hence R is a Sylow p -subgroup of E . It is clear also that R is a Sylow p -subgroup of G and $(|G : N_G(R)|, r) = 1$ for every prime $r \neq q$. Since $|\pi(G)| > 2$ by (c), R is normal in G . Hence G is ϕ -dispersive for some ordering ϕ of $\mathbb{P}$, a contradiction. The theorem is proved.

Corollary 4.3 (See [21, Theorem C]). *If every n -maximal subgroup of a soluble group G is $\mathcal{U}$ -subnormal in G and $|\pi(G)| \geq n$, then G is ϕ -dispersive for some ordering ϕ of the set of all primes.*

Finally, note that there are examples which show that the restrictions on $|\pi(G)|$ in Theorems A, B, and D cannot be weakened.

References

- [1] K. Doerk, T. Hawkes, Finite Soluble Groups, Walter de Gruyter, Berlin-New York, 1992.
- [2] A.N. Skiba, A Characterization of Finite Soluble Groups of Given Nilpotent Length, in: Problems in Algebra, Minsk, University Press 3 (1987) 21–31.
- [3] L.A. Shemetkov, A.N. Skiba, Formations of Algebraic Systems, Nauka, Moscow, 1989.
- [4] B. Huppert, Normalteiler und Maximal Untergruppen Endlicher Gruppen, Math. Z. 60 (1954) 409–434.

- [5] X.Y. Guo, K.P. Shum, Cover-avoidance properties and the structure of finite groups, *J. Pure Appl. Algebra* 181 (2003) 297–308.
- [6] W. Guo, K.P. Shum, A.N. Skiba, X -Semipermutable subgroups of finite groups, *J. Algebra* 315 (2007) 31–41.
- [7] Baojun Li, A.N. Skiba, New characterizations of finite supersoluble groups, *Sci. China Ser. A: Math.* 50 (1) (2008) 827–841.
- [8] W. Guo, A.N. Skiba, Finite groups with given s -embedded and n -embedded subgroups, *J. Algebra* 321 (2009) 2843–2860.
- [9] Shirong Li, Finite non-nilpotent groups all of whose second maximal subgroups are TI -groups, *Math. Proc. of the Royal Irish Academy* 100A (1) (2000) 65–71.
- [10] W. Guo, H.V. Legchekova, A.N. Skiba, Finite groups in which every 3-maximal subgroup permutes with all maximal subgroups, *Math. Notes* 86 (3) (2009) 325–332.
- [11] W. Guo, Yu.V. Lutsenko, A.N. Skiba, On nonnilpotent groups with every two 3-maximal subgroups permutable, *Siberian Math. J.* 50 (6) (2009) 988–997.
- [12] Yu.V. Lutsenko, A.N. Skiba, Structure of finite groups with S -quasinormal third maximal subgroups, *Ukrainian Math. J.* 61 (12) (2009) 1915–1922.
- [13] Yu.V. Lutsenko, A.N. Skiba, Finite groups with subnormal second and third maximal subgroups, *Math. Notes* 91 (5) (2012) 680–688.
- [14] A. Mann, Finite groups whose n -maximal subgroups are subnormal, *Trans. Amer. Math. Soc.* 132 (1968) 395–409.
- [15] A. Ballester-Bolinches, L.M. Ezquerro, *Classes of Finite Groups*, Springer-Verlag, Dordrecht, 2006.
- [16] L.A. Shemetkov, *Formations of Finite Groups*, Nauka, Moscow, 1978.
- [17] W. Guo, A.N. Skiba, On $\mathcal{F}\Phi^*$ -hypercentral subgroups of finite groups, *J. of Algebra* 372 (2012) 275–292.
- [18] A.N. Skiba, *Algebra of formation*, Belaruskaya Navuka, Minsk, 1997.
- [19] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [20] Otto-Uwe Kramer, Endliche Gruppen mit paarweise teilerfremden Indizes, *Math. Z.* 139 (1) (1974) 63–68.
- [21] V.A. Kovaleva, A.N. Skiba, Finite solvable groups with all n -maximal subgroups $\mathcal{U}$ -subnormal, *Sib. Math. J.* 54 (1) (2013) 65–73.

[22] D.J.S. Robinson, A course in the Theory of Groups, Springer, New York, 1982.